# SRILANKA'S HEALTHIEST WORKPLACE -AIA Wality

THE 2019 MARKET HEALTH REPORT FOR SRI LANKA



## TABLE OF CONTENTS

EXECUTIVE SUMMARY	4
HOW TO READ THIS REPORT	4
PART ONE - YOUR EMPLOYEES	6
OVERALL HEALTH RISK PROFILE	7
LIFESTYLE HEALTH	9
PHYSICAL ACTIVITY	10
NUTRITION	11
SMOKING	13
ALCOHOL	15
HEALTH PERCEPTION AND MOTIVATION TO CHANGE	16
SLEEP	18
MENTAL WELLBEING	20
WORK-RELATED STRESS	20
DEPRESSIVE SYMPTOMS	22
FINANCIAL CONCERNS	23
CLINICAL HEALTH	24
BODY MASS INDEX (BMI)	24
HEALTH SCREENING AND CLINICAL INDICATORS	25
MUSCULOSKELETAL CONDITIONS	26
CHRONIC CONDITIONS	26
ABSENCE, PRESENTEEISM AND PRODUCTIVITY	28
WORK ENGAGEMENT	30
JOB SATISFACTION AND WORK ENGAGEMENT	30
PART TWO - THE WORK ENVIRONMENT	33
LEADERSHIP AND CULTURE	34
ORGANISATIONAL SUPPORT	35
WORKPLACE HEALTH INTERVENTIONS - FACILITIES AND SERVICES	37
PHYSICAL ACTIVITY	37
NUTRITION	38
SMOKING	38
ALCOHOL	39
SLEEP	39
FATIGUE	39

MENTAL HEALTH AND WELLBEING	40
DIGITAL HEALTH	41
OTHER HEALTH SUPPORT	42
EVIDENCE FOR EMPLOYERS	43
SERVICES	43
INCENTIVES	44
PROMOTION	44
LEADERSHIP	45
EVALUATION	45
CONCLUSION	46
APPENDICES	47
APPENDIX A	48
DEMOGRAPHIC BREAKDOWN OF RESPONDENTS	48
APPENDIX B	50
YEAR-ON-YEAR RESULTS*	50
APPENDIX C	51
CALCULATION OF PRODUCTIVITY LOSS	51
APPENDIX D	53
RISK FACTOR DEFINITIONS USED IN THIS REPORT	53
APPENDIX E	54
THE TEAM BEHIND THE HEALTHIEST WORKPLACE COMPETITION	1 54
ABOUT AIA & THE HEALTHIEST WORKPLACE BY AIA VITALITY	54
RAND EUROPE	55
REFERENCES	56



#### **EXECUTIVE SUMMARY**

Thank you for taking part in the 2019 Sri Lanka's Healthiest Workplace by AIA Vitality. This year, for the second round of the survey, 47 organisations, representing a combined workforce of 2,137 employees, participated in the survey. The wider Healthiest Workplace by AIA Vitality survey, encompassing Australia, Hong Kong, Malaysia, Sri Lanka and Thailand, saw participation from 426 organisations, representing a combined workforce of 26,456 employees.

A growing body of research on corporate wellness indicates that proactive management of employees' physical and mental health can produce a range of important business benefits - including a reduction in absence, greater staff engagement and productivity, and a reduction in staff turnover. The aim of the Healthiest Workplace by AIA Vitality study is to understand how the workplace can affect employees' health, and further relate the health of employees to their performance and engagement at work. This should assist employers in building the case to invest in the health and wellbeing of their staff.

The study was developed by AIA Group Corporate Solutions (GCS) and is delivered in partnership with RAND Europe. Having inherited Britain's Healthiest Workplace that started in 2013, AIA brought this to Asia Pacific for the first time in 2017 to support organisations to improve understanding of their employees' personal health and wellbeing.

#### **HOW TO READ THIS REPORT**

This report analyses survey responses from all 47 organisations and 2,137 employees in Sri Lanka. Their results are benchmarked against other markets participating in the Healthiest Workplace by AIA Vitality, specifically Australia, Hong Kong, Malaysia and Thailand (combined in the Asia Pacific benchmark) and against Sri Lanka's 2018 results.

Figures presented in this report are rounded. Therefore percentages in graphs might not always add up to 100%.

The main outcomes and indicators from the survey are presented in the dashboard shown on the next page. A detailed analysis of the employees who took part in the survey is provided in Appendix A and your results compared to previous years can be found in Appendix B.

#### SRI LANKA'S DASHBOARD

CATEGORY	INDICATOR	SRI LANKA <sup>2</sup>	ASIA PACIFIC <sup>3</sup>	YOUR 2018 RESULTS⁴
Key outcomes				
Long-term health	AIA Vitality Age gap (in years) <sup>5</sup>	4.9	5.2	5.2
Short-term work impairment	Days lost per employee per year <sup>6</sup>	47.9	69.2	49.3
Work engagement	% low engaged <sup>7</sup>	10.3%	14.9%	8.3%
Determinants of h	ealth, productivity and engagement			
Lifestyle health	% with insufficient physical activity <sup>8</sup>	58.6%	60.5%	-
	% not eating a healthy diet	88.9%	85.6%	91.5%
	% current smokers	13.6%	8.1%	12.8%
	% exceeding alcohol guidelines	1.9%	1.7%	1.1%
	% sleeping less than 7 hours per night	46.4%	49.0%	46.4%
Clinical health	% obese	10.3%	14.0%	10.6%
	% with severe musculoskeletal problems <sup>9</sup>	7.4%	8.1%	7.6%
Mental health	% with moderate or severe symptoms of depression	15.0%	6.8%	12.9%
	% with 1 or more work-related stress factors	56.1%	49.8%	59.5%
	% with a lot of financial concerns	30.9%	21.5%	33.6%
Employee charact	eristics			
Survey respondent		59.1%	38.4%	61.3%
profile	Average age	32.0	35.6	32.5
	Average income	LKR 104,761	-	LKR 92,595

For full descriptions of these outcomes and determinants please see the relevant sections of the report - p. 9 for lifestyle health; p. 20 for mental welbeing; p. 24 for clinical health.

The green/amber/red colour codes correspond to significantly better/similar/significantly worse results than the Asia Pacific benchmark, defined as being 5% (for Vitality Age gap and days lost) or 2 percentage points better/worse than the benchmark, respectively. Based on 2,137 employee responses from 47 organisations from Sri Lanka. Based on 24,319 employee responses from 406 organisations from the Asia Pacific benchmark. Based on 2,221 employee responses from 53 organisations from Sri Lanka's 2018 results. AIA Vitality Age as described on p. 7. The AIA Vitality Age gap refers to the average difference between the AIA Vitality Age and actual age of the employees. Work impairment due to absence and presenteeism, as measured using the Work Productivity and Activity Impairment (General Health) scale, see Appendix C. Utrecht Work Engagement scale, see p. 30. The physical inactivity indicator in 2018 is not comparable to other years due to changes in the measurement method. 1

<sup>2</sup> 

<sup>3</sup> 

<sup>5</sup> 

<sup>6</sup> 

<sup>8</sup> measurement method.

Employees who had to take time off work in the past 12 months due to musculoskeletal pain or 9 discomfort. This question was not asked in Australia in 2018.



#### OVERALL HEALTH RISK PROFILE

Increasingly, research is being conducted into the impact of people's lifestyle choices on their health, mortality and productivity. The Global Burden of Disease Study 2017 revealed that non-communicable causes account for 80% of all global deaths [1], while research conducted using Sri Lanka's Healthiest Workplace by AIA Vitality data has shown that lifestyle behaviours are significant drivers of short-term productivity loss. As such, supporting employees to make healthier choices is a critical part of any human resources and productivity management strategy.

#### AIA VITALITY AGE

AlA Vitality Age was developed by Discovery Health Ltd in association with the University of Cape Town and is underpinned by hundreds of international evidence-based academic research studies.

AlA Vitality Age shows the impact of a range of lifestyle, clinical and mental wellbeing risks on an individual's long-term health. The difference between a person's AlA Vitality Age and their actual age is referred to as their AlA Vitality Age gap, which corresponds to the predicted impact of these risks on their life expectancy. If a participant is particularly fit and healthy, their AlA Vitality Age could be lower than their actual age. However, for the majority of people, AlA Vitality Age will be higher than their actual age.

From an employer's perspective, the lower the average difference between AIA Vitality Age and actual age, the healthier the participating employees in the workforce.


A strong link has been identified between the AIA Vitality Age gap and productivity loss. Across all participants in Sri Lanka's Healthiest Workplace by AIA Vitality, the results demonstrate that each additional year of AIA Vitality Age gap corresponds, on average, to 3 additional days of lost productive time per employee per year due to ill-health-related absenteeism and presenteeism.

For Sri Lanka, the average difference between respondents' AIA Vitality Age gap of 4.9 years and the benchmark averages of 5.2 years for the Asia Pacific and 5.2 years for Sri Lanka's 2018 results means that each of the employees could be gaining 0.9 and gaining 0.7 additional days of productive time per annum compared to the benchmarks, respectively.

AIA VITALITY AGE
IS BASED ON
MODIFIABLE LIFESTYLE
AND CLINICAL RISK
FACTORS.


98% OF YOUR
EMPLOYEES HAVE
AN AIA VITALITY
AGE HIGHER THAN
THEIR ACTUAL AGE,
COMPARED TO THE
BENCHMARKS OF 95%
AND 98%.

#### VITALITY AGE GAP OVERVIEW



- Vitality Age lower than or equal to actual age
- Vitality Age higher than actual age

#### DISTRIBUTION OF THE AIA VITALITY AGE GAP



#### LIFESTYLE HEALTH

Smoking, excessive alcohol consumption, poor nutrition and physical inactivity are important modifiable risk factors. Encouraging employees to make healthy choices will support the prevention of chronic diseases. It should also have a beneficial impact on productivity, work performance and engagement.

The following sections show the employees' behaviour with regard to lifestyle choices, highlight where they are most at risk, and offer some suggestions on how to positively influence this. More detail on workplace health interventions can be found on page 37.

For each risk factor, the following definition of risk applies.

**Physical activity risk** refers to the percentage of employees who are doing less than the recommended 150 minutes of moderate-intensity or 75 min of vigorous-intensity exercise each week.


**Nutrition risk** refers to the percentage of employees who indicate they do not eat a balanced diet. A balanced diet is one based on recommended servings of fruits, vegetables, and whole grains, avoiding excessive portions of high-fat meats, high-fat dairy, added fat, trans fats, salt and sugary drinks.

**Smoking risk** refers to the percentage of employees who currently smoke cigarettes, cigars or pipes.

**Alcohol risk** refers to the percentage of employees drinking more than the recommended limit of 14 units of alcohol each week.

The graph below illustrates the proportion of the employees who are at risk as a result of their lifestyle indicators being outside of the healthy range, compared to both the benchmarks.<sup>10</sup>

#### LIFESTYLE RISK INDICTORS




THE EMPLOYEES IN SRI LANKA ARE MOST AT RISK FOR NUTRITION.

<sup>10</sup> Physical inactivity in 2018 is not comparable to other years due to changes in the measurement method.



# YOUR EMPLOYEES' PHYSICAL ACTIVITY LEVELS



Physical activity per week


- High risk (<120 minutes)
- Medium risk (120-150 minutes)
- Low risk (151+ minutes)

#### PHYSICAL ACTIVITY

Physical activity guidelines recommend that people should do at least 150 minutes of moderate-intensity physical activity – five sessions of 30 minutes each per week [2], or 75 minutes of vigorous-intensity physical activity per week. However, research shows that even doing 120 minutes of at least moderate-intensity physical activity per week may reduce the risk of type 2 diabetes or metabolic syndrome [3].

The following graph compares the proportion of the employees who are not engaging in adequate amounts of physical activity to the Asia Pacific benchmark.<sup>11</sup> 59% of your employees are at risk, with physical activity levels below 150 minutes of moderate-intensity activity per week and below 75 minutes of vigorous activity, compared to 61% in Asia Pacific.

#### OVERALL PHYSICAL ACTIVITY RISK PROFILE



Most organisations indicate that their employees are sedentary for a large part of their working day. Sedentary time, defined as time spent sitting at desks or in meetings, also increases a person's risk for disease, beyond the negative impact of insufficient physical activity. As an employer, you may be able to take steps to encourage employees to be as active as possible during their time at work. This may include encouraging employees to:

- take the stairs instead of the lift
- walk, run or cycle during lunch
- walk, run or cycle part or all of their way to work
- walk to colleagues' desks instead of phoning or emailing them

Physical inactivity in 2018 is not comparable to other years due to changes in the measurement

Wearable devices and mobile apps are becoming increasingly popular to keep track of daily physical activities. Such devices can indicate, for example, the number of steps taken in a day or the number of calories burnt. Of the analysed employees, 24% indicate that they use a wearable health device such as a fitness band or a fitness activity tracker.

On average, organisations in Sri Lanka offer 2 of the 15 physical activity interventions asked about in the survey. A summary of the use and effectiveness of these interventions is provided in the table below, with a detailed analysis of each intervention provided in the section titled "Workplace health interventions - facilities and services", commencing on page 37.

#### EFFECTIVENESS OF PHYSICAL ACTIVITY INTERVENTIONS

	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS
% of participants aware of the interventions on offer	14.8%	21.0%	8.4%
% of participants participating in at least one of the interventions (of those who are aware)	61.6%	69.4%	61.6%
% of participants who feel the interventions positively impacted their health	90.8%	94.3%	85.1%

#### **NUTRITION**

One can assess an individual's diet by measuring the intake of certain types of food either positively or negatively associated with chronic diseases. These food groups are principally:

**Fruit and vegetables:** the consumption of fruit and vegetables is measured by the number of servings a person consumes in a day. The World Health Organization (WHO) and the Food and Agriculture Organization of the United Nations (FAO) have recommended that individuals consume at least 400 grams of fruit and vegetables a day [4], which is equivalent to five servings [5].

**Whole grains:** found in foods including wholegrain bread, cereals such as oats, barley, millet, brown rice and whole wheat pasta. The FAO recommends that carbohydrate foods, under which whole grains are listed, make up around 55% of the energy we eat [6].


**Added salt:** the WHO advises that adults eat no more than 5g of salt per day. As our diets are generally already high in salt, adding salt can constitute a health risk [7].

WITHIN THE KEY
NUTRITION ELEMENTS,
EMPLOYEES' DIETS ARE
POOREST IN THE AREA
OF WHOLE GRAINS.




The healthy range intakes of each of these food groups are indicated in Appendix D. Nutrition risk, as shown in the graph below, is assessed as a composite score on the basis of each of these different nutrition elements.

#### OVERALL NUTRITION RISK PROFILE




#### FRUIT AND VEGETABLE INTAKE



Overall, 100% of the employees are at risk for at least one nutritional element. Within the key nutrition elements, the employees' diets are poorest in the area of whole grains. In addition, in the side chart, we examine the severity of the employees' nutrition risk, using consumption of fruit and vegetables as an indicator of overall diet. Data collected through Sri Lanka's Healthiest Workplace by AIA Vitality has shown that fruit and vegetable intake is the nutritional element which can be most easily addressed in the workplace, with 32% of organisations across the whole Healthiest Workplace by AIA Vitality study indicating that they offer fresh fruit and vegetables.

#### EMPLOYEES AT RISK FOR KEY NUTRITION ELEMENTS



One of the barriers preventing people from eating healthily is that healthy food can be more expensive than unhealthy food. Subsidising healthier food items available in the workplace (often accompanied by increasing the cost of unhealthy foods) can be an effective way to help to overcome this barrier [8].

On average, organisations in Sri Lanka offer 3 of the 11 nutrition interventions asked about in the survey. A summary of the use and effectiveness of these interventions is provided in the table below, with a detailed analysis of each intervention provided in the section titled "Workplace health interventions - facilities and services", on page 38.

#### EFFECTIVENESS OF NUTRITION INTERVENTIONS

	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS
% of participants aware of the interventions on offer	17.2%	24.6%	11.2%
% of participants participating in at least one of the interventions (of those who are aware)	91.8%	91.9%	88.1%
% of participants who feel the interventions positively impacted their health	87.5%	90.2%	88.3%

#### **SMOKING**

The harmful effects of smoking are well known. Smoking, both active and passive, is associated with a range of cancers, coronary heart disease, heart attack, stroke, peripheral vascular disease (damaged blood vessels), cerebrovascular disease (damaged arteries that supply blood to your brain), and chronic bronchitis. [9]. The WHO estimated that tobacco consumption leads to approximately 8 million deaths per year globally [10]. The prevalence of daily smokers in Sri Lanka in 2017 was approximately 10% [11].

14% of employees are current smokers, compared to the average of 8% in the Asia Pacific and 13% in Sri Lanka's 2018 results.


10% of employees do not currently smoke, but have done so in the past. Of this group, 1% stopped smoking 15 or more years ago, meaning their risk of smoking-related diseases is significantly reduced. 8%, on the other hand, stopped smoking 5 years ago or less.

14% OF EMPLOYEES
ARE CURRENT
SMOKERS.

As well as the detrimental effects on employee health outcomes, the costs due to productivity loss contribute to the global economic cost of smoking. Combined, productivity loss and health care expenditure amount to 1.8% of the worldwide GDP [12]. Other indirect costs include the productivity lost to smoking breaks and the cost of cigarette-related fire damage [13].

The graphs below compare the proportion of employees who currently smoke to the various benchmarks, as well as providing a detailed breakdown of the number of cigarettes smoked each day by current smokers.


#### OVERALL SMOKING RISK PROFILE



On average, organisations in Sri Lanka offer less than 1 of the 6 smoking interventions asked about in the survey. A summary of the use and effectiveness of these interventions is provided in the table below, with a detailed analysis of each intervention provided in the section titled "Workplace health interventions - facilities and services", on page 38.

#### EFFECTIVENESS OF SMOKING INTERVENTIONS

SRI LANKA ASIA PACIFI % of participants aware of the interventions on offer 4.0% 15.0%	
interventions on offer  4.0%  15.0%  % of participants participating in	IC LK 2018 RESULTS
	1.6%
at least one of the interventions 33.3% 45.7% (of those who are aware)	57.7%
% of participants who feel the interventions positively impacted 89.3% 93.2% their health	80.0%




Percentage of respondents
by cigarettes per day

■ 21+ ■ 11-20 ■ 6-10 ■ 3-5 ■ 1-2

#### **ALCOHOL**

Akin to smoking, the negative effects of alcohol on health are well known. Sustained consumption of levels of alcohol above recommended guidelines have been associated with higher risks of mortality [14, 15]. Furthermore, both higher-than-average alcohol consumption and heavy episodic drinking have been linked to various cancers, heart disease and stroke [16]. Given the increasing evidence of the adverse health effects of alcohol, the WHO launched a global strategy to reduce the harmful effects of alcohol [17]. In addition, in some countries consuming alcohol is not encouraged or is prohibited for Muslims (e.g. in Malaysia).


#### OVERALL ALCOHOL RISK PROFILE



2% of employees exceed recommended weekly consumption limits; 10% binge drink at least once a month.


On average, organisations in Sri Lanka offer less than 1 of the 3 alcohol interventions asked about in the survey. A summary of the use and effectiveness of these interventions is provided in the table below, with a detailed analysis of each intervention provided in the section titled "Workplace health interventions - facilities and services", on page 39.

#### **HEAVY DRINKERS:** DRINKING PATTERNS





#### **BINGE DRINKERS**



Frequency of binge drinking



#### EFFECTIVENESS OF ALCOHOL INTERVENTIONS

	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS
% of participants aware of the interventions on offer	4.9%	10.7%	3.7%
% of participants participating in at least one of the interventions (of those who are aware)	31.3%	48.2%	57.9%
% of participants who feel the interventions positively impacted their health	45.8%	95.9%	100.0%

#### HEALTH PERCEPTION AND MOTIVATION TO CHANGE

People are less likely to make changes to their lifestyles if they perceive themselves to be healthy. Therefore, it is important for employees to have a realistic perception of their health. Sometimes employees with multiple risk factors perceive themselves to be in good or very good health. Though some of the risk factors presented may be mild, not symptomatic, or controlled, perceived good health may indicate specific cases in which employees do not have good awareness of underlying risks to their health.

The graph below shows how your employees perceive their health relative to the number of risk factors<sup>12</sup> they actually have.

#### EMPLOYEES' HEALTH PERCEPTION AND RISK PROFILE




12% OF EMPLOYEES
HAVE FOUR OR MORE
RISK FACTORS, 56% OF
WHICH BELIEVE THEY
ARE IN 'GOOD' OR
'VERY GOOD' HEALTH.

<sup>12</sup> These include high BMI, low physical activity, poor nutrition, current smoking, and excessive alcohol consumption.

Awareness of our risk factors is not always sufficient to actually improve our health. Motivation is considered a key component of behavioural change and understanding employees' motivations can be crucial to designing effective interventions [18]. The areas in which respondents in Sri Lanka are most motivated to change are physical activity and smoking.

#### MOTIVATION TO CHANGE BEHAVIOUR BY RISK FACTOR








46% OF EMPLOYEES REPORT GETTING LESS THAN 7 HOURS OF SLEEP PER NIGHT.


#### **SLEEP PATTERNS**



#### SLEEP

Long working hours, increasing out-of-hours connectivity and work-related stress are all factors associated with sleep problems in workers who would not normally be classified as being at risk for sleep disorders. Furthermore, sleep problems have been shown to increase the probability of developing severe mental health conditions [19]. In addition, workers who report shorter amounts of sleep are more likely to avoid social interactions at work [20]. The graphs below show the hours of sleep reported by employees. 46% of employees reported getting less than 7 hours sleep in a 24-hour period, the optimal level of sleep from both a health and productivity perspective.


#### SLEEP PATTERNS: HOURS OF SLEEP PER NIGHT



Employees sleeping less than 7 hours per night

Poor sleep quality is associated with impairments at work, such as reduced concentration and difficulties with organising work. Among the analysed respondents, 13% indicated they had poor or very poor quality sleep during the week preceding the survey. The following graph shows the issues mentioned by the employees which are preventing them from sleeping well.

#### SLEEP QUALITY PATTERNS




On average, organisations in Sri Lanka offer 1 of the 5 sleep interventions asked about in the survey. A summary of the use and effectiveness of these interventions is provided in the table below, with a detailed analysis of each intervention provided in the section titled "Workplace health interventions - facilities and services", on page 39.

#### **EFFECTIVENESS OF SLEEP INTERVENTIONS**

	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS
% of participants aware of the interventions on offer	6.8%	12.9%	6.2%
% of participants participating in at least one of the interventions (of those who are aware)	73.3%	71.0%	65.0%
% of participants who feel the interventions positively impacted their health	86.1%	93.7%	93.1%

#### SLEEP QUALITY





#### MENTAL WELLBEING

Mental health and wellbeing is related to a wide range of factors such as poor lifestyle behaviours, health concerns, and financial concerns. Other workplace issues such as stress, organisational culture, leadership, and bullying can also impact upon the mental health and wellbeing of staff. Increasingly, mental health is being recognised as one of the key drivers of general health and wellbeing, and has been found to substantially impact productivity and engagement in the workplace.

#### **WORK-RELATED STRESS**

Work-related stress represents a significant problem internationally, and can affect organisational performance, including absenteeism, productivity and turnover. In the UK for example, work-related stress accounts for around two-thirds of all new cases of work-related ill-health. It is estimated that about 10.5 million working days are lost yearly due to work-related stress, depression and anxiety [21]. High levels of work-related stress can impact staff absence and turnover and negatively affect the productivity and performance levels of employees.

In Sri Lanka's Healthiest Workplace by AIA Vitality, work-related stress was measured using seven items from the UK Health & Safety Executive (HSE) Management Standards Indicator Tool.<sup>13</sup> Using this tool, stress is measured using the following statements:

**Control:** I have a choice in deciding what I do at work (never/seldom)

**Role:** I am clear what my duties and responsibilities are (never/seldom)

**Managerial support:** My line manager encourages me at work (disagree/strongly disagree)

**Peer support:** I receive the respect at work I deserve from my colleagues (disagree/strongly disagree)


**Change:** Staff are always consulted about change at work (disagree/strongly disagree)

**Relationship:** Relationships at work are strained (agree/strongly agree); and, I am subject to bullying at work (often/always)

**Demands:** I have unrealistic time pressures (often/always)

<sup>3</sup> http://www.hse.gov.uk/stress/standards/notesindicatortool.htm


#### **OVERALL STRESS RISK PROFILE**



Employees at risk for at least one work-related stress factor


56% of the analysed employees are subject to at least one dimension of work-related stress. Additionally, 26% of employees were subject to bullying at the workplace, 5% of whom indicated that they were bullied often or always.

#### BREAKDOWN OF WORK-RELATED STRESS INDICATORS



15% OF EMPLOYEES
HAVE MODERATE
TO HIGH LEVELS
OF ANXIETY OR
DEPRESSIVE
SYMPTOMS.

# SEVERITY OF DEPRESSIVE SYMPTOMS




#### **DEPRESSIVE SYMPTOMS**

The Kessler Psychological Distress Scale [22], a six-item questionnaire, is intended to yield a measure of distress based on questions about anxiety and depressive symptoms that a person has experienced in the most recent four-week period.

Depression, anxiety and stress can impact significantly on both mental and physical health. Stress management techniques and, where necessary, professional counselling can help address this risk factor in the work context [23].

#### OVERALL ANXIETY AND DEPRESSION RISK PROFILE



2...projeco m.a. moderate to mgn doprecente symptoms

On average, organisations in Sri Lanka offer 3 of the 14 mental wellbeing interventions asked about in the survey. A summary of the use and effectiveness of these interventions is provided in the table below, with a detailed analysis of each intervention provided in the section titled "Workplace health interventions - facilities and services", on page 40.


#### EFFECTIVENESS OF MENTAL HEALTH AND WELLBEING INTERVENTIONS

	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS
% of participants aware of the interventions on offer	13.2%	18.0%	9.1%
% of participants participating in at least one of the interventions (of those who are aware)	59.5%	59.8%	55.7%
% of participants who feel the interventions positively impacted their health	78.1%	92.2%	88.0%


#### **FINANCIAL CONCERNS**

One factor potentially underlying mental health problems is financial concerns, which can cause depression and damage relationships [24]. It is measured using the following question: "Do you have financial concerns at present?". 31% of the analysed employees noted that they had a lot of financial concerns at present.

#### OVERALL FINANCIAL CONCERNS RISK PROFILE



# EXTENT OF FINANCIAL CONCERNS






#### CLINICAL HEALTH

Clinical risk factors can refer to either a range of indicators that act as precursors for disease, or to existing health conditions. This report classifies these as body mass index, biometric indicators such as blood pressure and cholesterol, musculoskeletal conditions, and chronic health conditions. Many of these clinical factors are closely linked to the lifestyle behaviours outlined previously.

#### **BODY MASS INDEX (BMI)**

BMI is a measure of a person's weight in relation to their height. A healthy BMI is considered to be in the range 18.5 to 24.9. A BMI of 25 or over puts a person at increased risk for heart disease, high blood pressure, and even certain cancers [25, 26, 27]. Employees with a BMI of 30 or more or of less than 18.5 are classified as obese and underweight, respectively. Studies show that even modest weight loss (around 5%) achieved by people who are obese can have beneficial effects on health, medical expenditures, and improve quality of life [28].

#### OVERALL BODY MASS INDEX RISK PROFILE



41% of the analysed employees are either overweight or obese. Of the employees who have a BMI classified as 'overweight', 94% reported that their waist circumference is within the healthy range (less than 88 cm for women and less than 101cm for men) and are therefore classified as low

risk. In particular, waist circumference can be a more accurate measure of body composition for those with large muscle mass. Adjusting for waist circumference, the percentage of the employees who are overweight or obese reduces to 12%, compared to a the benchmarks of 17% for the Asia Pacific and 14% for Sri Lanka's 2018 results.

#### HEALTH SCREENING AND CLINICAL INDICATORS

Employees can monitor their clinical risk factors by undergoing regular screening tests to assist in the early detection and treatment of chronic diseases.

The three most commonly screened biometric indicators are blood pressure, cholesterol and blood glucose.

**High blood pressure,** even 'minor high' blood pressure (referred to as prehypertension), is a risk factor for a range of conditions - in particular, stroke [29]. Values of 140 over 90 are considered as first stage hypertension according to international standards, with values above that posing increasing health risk [30].

**Cholesterol** is measured in milligrammes per decilitre of blood (mg/dL). While cholesterol is a product produced by the body, it is also found in foods, which can raise cholesterol in the body to unhealthy levels.

**Blood glucose,** or blood sugar, can be measured in different ways, and the survey asked about non-fasting or random blood sugar levels.

33% of the analysed employees have had all three of blood pressure, cholesterol and glucose tested in the past 12 months.

The tables below summarise both employees' awareness of, and the percentage who are at elevated risk for, these clinical factors.

HEALTH SCREENING PARTICIPATION AND RISK


	PERCENTAGE OF RESPONDENTS TESTED IN PAST 12 MONTHS (PERCENTAGE AT RISK IN PARENTHESES)*			
	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS	
Blood pressure	64% (47%)	83% (42%)	68% (45%)	
Cholesterol	40% (8%)	56% (9%)	46% (6%)	
Blood glucose	46% (3%)	56% (3%)	50% (4%)	
All 3	33% (0%)	47% (0%)	37% (0%)	

<sup>\*</sup> Of those who recalled their result.

33% OF EMPLOYEES
HAVE TESTED FOR
THEIR FULL SET
OF BIOMETRIC
INDICATORS IN THE
LAST 12 MONTHS:
BLOOD PRESSURE,
CHOLESTEROL AND
GLUCOSE

75% OF EMPLOYEES REPORT SYMPTOMS OF ONE OR MORE MUSCULOSKELETAL CONDITIONS.

# COMMON MUSCULOSKELETAL SYMPTOMS




#### MUSCULOSKELETAL CONDITIONS

Musculoskeletal (MSK) conditions relate to any health problem involving muscles, bones and joints. MSK problems are very common, and many forms may arise through contemporary forms of office work. Indeed, lack of positive practice to support people with musculoskeletal disorders in work can have different types of costs, from lost earnings, to reduced productive working time and early retirement, with an accompanying strain on both household incomes and the national welfare system [31, 32].

75% of employees reported one or more musculoskeletal conditions, compared to the benchmarks of 82% and 72% for the Asia Pacific and Sri Lanka's 2018 results, respectively.

The most common symptoms reported in Sri Lanka are lower back and neck while 25% of the employees report no symptoms. 22% of the analysed employees indicate that their condition prevented them from doing normal work at home or away from home and 7% indicate that the pain or discomfort caused them to take time off work.

#### OVERALL MUSCULOSKELETAL CONDITIONS RISK PROFILE



#### **CHRONIC CONDITIONS**

Chronic conditions have become much more common as populations age and new treatments allow people to live longer with diseases. There are a number of important lifestyle factors that increase the risk for developing a chronic condition, such as smoking, poor diet, lack of exercise, and

harmful alcohol use, among others. Employees reported whether they had been told by a doctor that they had one or more of the following chronic disease risk factors or conditions including kidney conditions, high blood pressure, high cholesterol, diabetes, heart disease, stroke or cancer. 30% of the analysed employees reported one or more chronic conditions compared to 30% in the Asia Pacific and 29% in Sri Lanka's 2018 results.

The table below shows the percentage of employees at risk for preventable common chronic conditions; in other words, those that may be prevented through early detection, improved diet, exercise, and treatment therapy.

#### EMPLOYEES AT RISK FOR PREVENTABLE COMMON CHRONIC CONDITIONS

	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS
Heart condition or disease	1.2%	1.3%	0.8%
Kidney condition or disease	0.7%	0.6%	1.3%
Cancer	0.3%	0.5%	0.0%
Diabetes	3.3%	2.6%	3.3%
High blood pressure	3.7%	7.3%	2.8%

On average, organisations in Sri Lanka offer 4 of the 19 health support interventions asked about in the survey. A summary of the use and effectiveness of these interventions is provided in the table below, with a detailed analysis of each intervention provided in the section titled "Workplace health interventions - facilities and services", on page 41.

#### FFFECTIVENESS OF HEALTH SUPPORT INTERVENTIONS

	SRI LANKA	ASIA PACIFIC	LK 2018 RESULTS
% of participants aware of the interventions on offer	17.6%	23.1%	13.1%
% of participants participating in at least one of the interventions (of those who are aware)	72.5%	81.0%	70.9%
% of participants who feel the interventions positively impacted their health	86.2%	90.8%	82.3%

ACCORDING TO
EMPLOYEES' SELFREPORTED ABSENCE
AND PRESENTEEISM,
ORGANISATIONS IN
AUSTRALIA ARE LOSING
AN AVERAGE OF 47.9
DAYS PER YEAR PER
EMPLOYEE.

### ABSENCE, PRESENTEEISM AND PRODUCTIVITY

There is substantial evidence concerning the relationship between employee health and wellbeing and an organisation's productivity. <sup>14</sup> The link between an organisation's productivity and employee health and wellbeing can be explored through two measures - absence and presenteeism.

Absence refers to the actual absence of employees from work. Absence due to health problems is relatively easy to measure as detailed records of employee sickness absence are generally kept by organisations.

Presenteeism due to health problems refers to reduced productivity at work. While present at work, employees are limited or constrained by health problems to carry out their daily activities, resulting in productivity loss [33]. Contrary to absence, presenteeism due to ill-health is more difficult to measure, but a range of survey tools are currently available to obtain a proxy of an organisation's presenteeism levels.<sup>15</sup>

One widely validated tool to measure self-reported absence and presenteeism – and in turn, productivity loss – is the Work Productivity and Activity Impairment (WPAI) Questionnaire. <sup>16</sup> The WPAI has been used in a wide range of studies to measure the loss of productivity from both general health problems as well as from specific conditions. In the Sri Lanka's Healthiest Workplace by AIA Vitality survey, the general health WPAI (WPAI-GH) has been used.


The graph overleaf expresses in days the lost productive time experienced by the analysed employees as a result of health-related absence and presenteeism. This is calculated using the employees' answers to the WPAI questionnaire and their self-reported working hours per week.

<sup>14</sup> For an introduction see [33].

For an overview of the methods used to measure presententeeism see [34, 35].

<sup>6</sup> All information on the WPAI can be found here: www.reillyassociates.net/WPAI\_General.html

#### OVERALL PRODUCTIVITY PROFILE



Expressed as a percentage of available working hours, the analysed employees lost 18.4% of working hours due to ill-health related absence (2.3%) and presenteeism (16.1%) in the week prior to the survey, compared to 26.6% in the Asia Pacific and 18.9% in Sri Lanka's 2018 results. This equates to 47.9 days lost per employee per year on average for Sri Lanka (assuming a five day working week). In order to quantify the productivity loss due to absence and presenteeism, the number of lost working hours is multiplied by the compensation (wage) that employees receive. This method of costing is referred to in the literature as the Human Capital Approach (HCA) and is widely used because of its computational ease.<sup>14</sup>

For Sri Lanka, the estimated average monthly cost of health-related absence and presenteeism per organisation is estimated at LKR 12,424,019 (1,783,198 due to absence and 10,640,822 due to presenteeism).

Please note that this estimate is extrapolated from the employees who completed the survey, and is based on their self-reported absence and presenteeism during the week prior to the survey. More detail and information on determining productivity loss can be found in Appendix C.

Other methods also exist. However, these require much more data collection in order to generate estimates.



#### **WORK FNGAGEMENT**

Research conducted using Britain's Healthiest Workplace data has demonstrated a clear relationship between ill-health and lost productivity [36]. Indeed, the analysis shows that distinct lifestyle behaviours, for example, exercise or nutrition habits, can impact significantly on short-term productivity through presenteeism. This loss is beyond the productive time loss that employers experience due to employee absence, often driven by chronic health conditions. Two other key factors influencing employee productivity are job satisfaction and work engagement, which are discussed in detail below.


#### JOB SATISFACTION AND WORK ENGAGEMENT

Employee engagement refers to the amount of energy, dedication and focus people bring to their work. It is currently regarded as one of the key 'people' factors that differentiate higher and lower levels of organisational performance on a range of indicators. Evidence suggests that there is a relationship between improved lifestyle and clinical health, and increased work engagement.

The graph overleaf indicates employees' satisfaction with their jobs. This in turn reflects on the levels of employee engagement. The assumption underlying this measure is that satisfied employees are engaged and full of energy, which in turn will contribute to organisational performance. The link between employee satisfaction and organisational performance has been explored in various studies, which found that employee satisfaction and engagement can impact business outcomes for a wide range of organisations [37].

Job satisfaction and work engagement are measured in a number of ways. A commonly asked question to measure job satisfaction is the level of agreement with the statement: "all in all I am satisfied with my job". The response to this question is shown in the graph overleaf.

#### BREAKDOWN OF EMPLOYEE JOB SATISFACTION



Work engagement comprises a wider set of dimensions related to an employee's attitude to their job and their organisation. Like job satisfaction, it is assumed to be the opposite of 'burn-out' [38]. To measure work engagement the Utrecht Work Engagement Scale-9 (UWES-9) was used, which captures the following three dimensions of work engagement through nine questions:

#### Vigour

- At my work, I feel bursting with energy
- At my job, I feel strong and vigorous
- When I get up in the morning, I feel like going to work

#### **Dedication**

- I am enthusiastic about my job
- I am proud of the work that I do
- My job inspires me


#### **Absorption**

- I am immersed in my work
- I get carried away when I am working
- I feel happy when I am working intensely

These three dimensions are defined as follows [38]: "Engagement is a positive, fulfilling, work-related state of mind that is characterised by vigour, dedication, and absorption. Rather than a momentary and specific state, engagement refers to a more persistent and pervasive affective-cognitive state that is not focused on any particular object, event, individual, or behaviour. Vigour is characterised by high levels of energy and mental resilience while working, the willingness to invest effort in one's work, and persistence even in the face of difficulties. Dedication refers to being strongly involved in one's work and experiencing a sense of significance, enthusiasm, inspiration, pride, and challenge. Absorption is characterised by being fully concentrated and happily engrossed in one's work, whereby time passes quickly and one has difficulties with detaching oneself from work."


Overall 18% of the employees scored high or very high for Work Engagement compared to the benchmarks of 18% in the Asia Pacific and 22% in Sri Lanka's 2018 results. The proportion of employees scoring low or very low on each of these dimensions is shown in the graph.

#### OVERALL WORK ENGAGEMENT PROFILE



Employees with low or very low levels of engagement

# PERCENTAGE OF LOW ENGAGED





There is a growing recognition for the need to develop a comprehensive organisational approach to align health promotion and human resource management practices. The quality of an employee's workplace experience has an impact on health and wellbeing. A direct relationship exists between job design, work-life balance, organisational change, and the health of the workforce. Work intensification and reorganisation, often coupled with technological change, has contributed to an increased incidence of musculoskeletal disorders and mental ill-health. In designing jobs and developing management practices, organisations are urged to make their staff feel valued by the organisation. This process motivates employees to deliver a quality product or service without excessive stress that might lead to poor health [39].



#### LEADERSHIP AND CULTURE

It is increasingly recognised that staff health and wellbeing is more than a matter of purely individual attention. The health, safety and wellbeing of staff can directly contribute to organisational success; a healthy workforce reduces costs in the form of sickness absence, lowers spending on staff turnover, reduces ill-health retirement and subsequently increases productivity [40]. Senior management, by taking an interest in health and wellbeing issues, are increasingly aware of the key elements to improving health and wellbeing at the workplace [41]. Line managers should play a key role in ensuring that workplaces are settings that promote good health. Good practice ensures that performance on staff health and wellbeing is regularly reported on, and discussed, at board or other management levels - and that action is taken on any existing and emerging problems with regard to the health and wellbeing of the workforce. Efficient practice further ensures that staff health and wellbeing is at the heart of line manager training, development and appraisal. At the same time, line managers should understand that the health and wellbeing of the workforce is their responsibility, and organisations should support them to act accordingly [41].


#### PERCEPTION OF LEADERSHIP AMONG THE EMPLOYEES

	% OF PARTICIPANTS WHO AGREE/STRONGLY AGREE		
	SRI LANKA	ASIA PACIFIC	SRI LANKA'S 2018 RESULTS
My line manager cares about my health and wellbeing	66%	57%	65%
My line manager encourages me at work	72%	61%	73%
Leaders view the level of employee health and wellbeing as an important indicator of the organisation's success.	58%	63%	60%

#### **ORGANISATIONAL SUPPORT**


Even in areas where the employees are less willing to change of their own volition, creating an environment conducive to change and a culture of health within the organisation may provide them with extra motivation [42]. The graph below shows the areas in which employees indicate that they feel supported by their organisation.

#### EMPLOYEE PERCEPTION OF ORGANISATIONAL SUPPORT



Linked to this, and as described earlier, the areas in which the greatest number of the employees are motivated to change are physical activity and smoking.

#### EMPLOYEE MOTIVATION TO CHANGE



<sup>\*</sup> The percentages for alcohol and smoking are based solely on those employees who consume alcohol or are active smokers.



Organisational support can express itself in various ways and the table below indicates how deeply health and wellness is embedded in an organisation's culture, through assessing levels of both investment and accountability for employee wellbeing.

#### KEY CULTURAL FACTORS

	PERCENTAGE/AVERAGE OF ORGANISATIONS		
	SRI LANKA	ASIA PACIFIC	SRI LANKA'S 2018 RESULTS
Budget for health and wellbeing facilities and services exists	57%	61%	65%
Staff health and wellbeing is discussed at the board level at least once a year	90%	85%	83%
Staff health and wellbeing is discussed at the executive committee level at least once a year	85%	87%	94%
Staff health and wellbeing is discussed at the middle management level at least once a year	94%	86%	91%
Staff health and wellbeing is discussed at all staff level at least once a year	97%	87%	84%
Providing incentives for participation and/or recognising or rewarding employees for healthy behaviour and health improvement	20%	42%	35%

# WORKPLACE HEALTH INTERVENTIONS – FACILITIES AND SERVICES

Expanding on the summarised information provided earlier, the following tables provide an overview of the interventions offered by organisations in Sri Lanka and the share of other organisations in the Healthiest Workplace by AIA Vitality (AIA HW) survey that offer those same interventions. In addition, the tables detail the percentage of employees aware, using and benefitting from these interventions, benchmarked against awareness, utilisation and perceived benefit amongst employees from other organisations that offer the interventions.

#### PHYSICAL ACTIVITY

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Provision of information on physical activity	25%	29%	11% (22%)	70% (67%)	98% (92%)
Onsite gym or fitness facility	22%	32%	34% (39%)	47% (61%)	90% (93%)
Offsite gym / health club membership discount	43%	29%	20% (21%)	48% (47%)	86% (93%)
Bicycle storage facilities	8%	19%	14% (19%)	53% (42%)	89% (96%)
Bicycle purchase scheme	3%	2%	7% (19%)	0% (64%)	n/a (95%)
Fitness classes	25%	34%	22% (28%)	43% (51%)	97% (93%)
Locker room with showers available at worksite	28%	27%	13% (25%)	57% (60%)	86% (92%)
Walk or cycle to work	13%	15%	15% (19%)	44% (68%)	91% (98%)
Sponsored walks or runs	28%	34%	13% (22%)	59% (61%)	87% (94%)
Corporate challenges	20%	35%	16% (18%)	58% (58%)	94% (92%)
Running clubs or other informal groups	17%	27%	14% (18%)	43% (57%)	86% (96%)
Bootcamps	0%	9%	n/a (26%)	n/a (65%)	n/a (97%)
Wearable fitness trackers or apps	10%	20%	8% (16%)	70% (77%)	95% (92%)
Stairs initiatives (e.g. inviting stairs or stair challenges)	28%	21%	12% (16%)	85% (69%)	82% (94%)
Other exercise opportunities (e.g. walking trails)	8%	13%	9% (6%)	57% (77%)	100% (98%)

## NUTRITION

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Healthy eating information	47%	36%	23% (32%)	76% (76%)	91% (93%)
Talks or workshops on nutrition and its health impacts	25%	34%	14% (23%)	67% (67%)	94% (92%)
Overweight and body fat assessment for customised nutritional advices	25%	29%	11% (19%)	68% (70%)	91% (95%)
Calorie intake calculator or canteen menus with calorie information	10%	14%	12% (21%)	70% (74%)	95% (94%)
Healthy food alternatives at canteens	40%	17%	18% (25%)	85% (81%)	89% (95%)
Healthy food alternatives in vending machines	10%	8%	7% (10%)	65% (40%)	80% (87%)
Fresh fruit and vegetables in the workplace	30%	32%	15% (29%)	88% (89%)	93% (92%)
Dietician/nutritionist services	17%	14%	11% (25%)	52% (62%)	94% (96%)
Access to fresh drinking water (other than tap water)	63%	59%	28% (28%)	96% (95%)	89% (90%)
Access to a microwave	65%	59%	27% (29%)	76% (79%)	72% (78%)
Access to a fridge	68%	59%	23% (30%)	86% (87%)	75% (80%)

## **SMOKING**

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Smoking cessation information	13%	21%	11% (23%)	34% (44%)	79% (90%)
Online smoking cessation programme	5%	2%	2% (20%)	0% (36%)	n/a (94%)
Group smoking cessation programme	3%	3%	3% (19%)	0% (34%)	n/a (94%)
Individual smoking cessation programme	0%	8%	n/a (12%)	n/a (29%)	n/a (87%)
Cognitive Behavioural Therapy or counselling	5%	9%	2% (15%)	100% (43%)	100% (95%)
Other assistance to quit smoking	5%	4%	2% (2%)	0% (48%)	n/a (100%)

## ALCOHOL

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Information on problem drinking	13%	13%	8% (21%)	27% (51%)	88% (95%)
Alcohol counselling	13%	7%	3% (10%)	20% (33%)	50% (93%)
Other assistance related to problem drinking	3%	3%	3% (1%)	100% (50%)	0% (100%)

## SLEEP

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Information on good sleeping habits	25%	20%	8% (16%)	62% (76%)	88% (94%)
Place where you can rest	38%	25%	17% (20%)	76% (70%)	84% (95%)
Apps/programmes promoting healthy sleep	5%	5%	1% (20%)	0% (76%)	n/a (89%)
Events promoting healthy sleep	5%	7%	2% (9%)	0% (70%)	n/a (97%)
Other	0%	2%	n/a (0%)	n/a (0%)	n/a (n/a)

## **FATIGUE**

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Information on fatigue	17%	15%	10% (17%)	59% (67%)	92% (90%)
Fatigue management interventions	8%	11%	2% (19%)	25% (63%)	100% (97%)

### MENTAL HEALTH AND WELLBEING

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Training on common mental health conditions (such as depression, anxiety disorders etc.)	28%	20%	22% (21%)	62% (49%)	76% (86%)
Resilience, energy or stress management classes or programmes	25%	20%	12% (17%)	45% (52%)	93% (89%)
Mindfulness classes or programmes	28%	19%	12% (24%)	61% (54%)	83% (91%)
Massage or relaxation classes or programmes	10%	13%	5% (23%)	70% (51%)	57% (91%)
Counselling or psychotherapy services	25%	17%	23% (19%)	24% (36%)	74% (94%)
Employee Assistance Programmes	38%	28%	19% (26%)	41% (30%)	78% (90%)
Workload or time management training	40%	23%	13% (16%)	58% (56%)	90% (92%)
Volunteering or charity work	52%	45%	18% (22%)	63% (62%)	87% (88%)
Coaching	43%	25%	14% (16%)	46% (51%)	85% (92%)
Other mental health support - onsite/telephone/mobile app/online	17%	13%	12% (14%)	26% (56%)	79% (96%)
Cognitive Behavioural Therapy or other types of psychological therapy	3%	6%	0% (14%)	n/a (45%)	n/a (99%)
Mental health and wellbeing information	25%	26%	18% (18%)	56% (54%)	74% (92%)
Financial wellbeing courses or programmes	17%	23%	10% (14%)	55% (59%)	91% (92%)
Other	5%	2%	6% (8%)	67% (96%)	50% (100%)

## DIGITAL HEALTH

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Wellbeing app targeting a broad range of physical health, mental health and lifestyle issues	17%	15%	12% (25%)	67% (82%)	89% (91%)
Wellbeing app targeting specific health issues, such as weight, exercise, or mental health	8%	14%	12% (20%)	71% (76%)	86% (91%)
Online coaching	5%	6%	9% (14%)	59% (59%)	70% (97%)
Surveys	3%	8%	3% (5%)	60% (77%)	67% (92%)
Digital platform incentivising healthy behaviours	13%	13%	3% (14%)	25% (71%)	100% (91%)
Other	0%	3%	n/a (13%)	n/a (99%)	n/a (99%)

### OTHER HEALTH SUPPORT

	% OF ORGANISATIONS IN SRI LANKA THAT OFFER THE INTERVENTION	% OF ORGANISATIONS IN AIA HW THAT OFFER THE INTERVENTION	% OF EMPLOYEES BEING AWARE OF THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THEY'VE USED THE INTERVENTION. (AIA HW BENCHMARK IN PARENTHESES)	% OF EMPLOYEES INDICATING THE INTERVENTION IMPROVED THEIR HEALTH. (AIA HW BENCHMARK IN PARENTHESES)
Provision of health information	50%	52%	32% (38%)	58% (68%)	81% (85%)
Clinical screening (e.g., blood glucose, blood pressure)	55%	59%	28% (45%)	66% (79%)	84% (87%)
Condition specific screening, such as for cancer	17%	26%	10% (18%)	25% (62%)	77% (93%)
Disease management (management of long term conditions such as diabetes, asthma, chronic obstructive pulmonary disease)	17%	23%	15% (20%)	50% (58%)	68% (93%)
Musculoskeletal disorders prevention	13%	20%	6% (17%)	18% (57%)	100% (94%)
Support in returning to work after illness	68%	35%	19% (17%)	51% (45%)	84% (93%)
Employee assistance programme	40%	35%	22% (28%)	47% (42%)	87% (93%)
Nurse advice line	13%	14%	25% (22%)	56% (57%)	85% (90%)
Occupational health / safety programme	40%	48%	33% (32%)	67% (65%)	90% (89%)
Vaccination (e.g. flu vaccine) onsite or in partnered clinics	5%	37%	2% (33%)	0% (71%)	n/a (95%)
On-site health clinics / medical services	32%	27%	26% (34%)	67% (81%)	84% (91%)
Wellness days where employees have health checks and get advice on improving their wellbeing	47%	42%	22% (28%)	64% (76%)	86% (89%)
Team discussions and activities	30%	35%	14% (16%)	81% (72%)	84% (91%)
Workshops on physical and mental health issues	38%	29%	24% (21%)	68% (62%)	75% (89%)
GP advice line	8%	14%	11% (20%)	50% (44%)	100% (90%)
Virtual GP service (telemedicine)	5%	5%	10% (9%)	69% (43%)	100% (89%)
Women health and wellness talks or workshops: fertility, healthy family planning, workplace and motherhood, disease and cancer prevention, travel medicine prevention/communicable disease prevention	22%	15%	16% (19%)	45% (56%)	83% (92%)
Men health and wellness talks or workshops: fertility, disease and cancer prevention, travel medicine prevention/communicable disease prevention	17%	11%	11% (12%)	36% (45%)	82% (88%)
Children disease prevention	5%	7%	9% (9%)	36% (42%)	100% (94%)

# EVIDENCE FOR EMPLOYERS

Workplace wellness programmes typically aim to change health-related behaviours of employees, such as getting smokers to quit, supporting weight loss, or promoting physical fitness. When studies assess the outcomes of health and wellbeing or wellness programmes, positive effects are found for about half of the studies [43]. Some studies have shown that mere participation in any workplace wellness programme can have an impact, as it can stimulate participants to adopt more healthy lifestyles [42]. The likelihood that programmes will actually achieve these outcomes depends on a number of factors. A recent study identified some key facilitators through a review of academic studies which may help in the design of workplace wellness programmes [44]. These can be captured as follows:

- 1. **Services** a programme needs to offer a good range of services of interest to employees and that are grounded in evidence
- 2. Incentives a programme needs to create opportunities for employees to engage. A variety of mechanisms such as ensuring access, and monetary and non-monetary rewards, can improve engagement
- **3. Promotion** broad outreach and various means of dissemination can help to inform employees of the existence and importance of a specific initiative or health and wellbeing programme
- **4. Leadership** support from both board level and direct supervisors is important to generate widespread participation
- **5. Evaluation** better programmes always aim to improve quality by monitoring progress and soliciting feedback from staff

We look at these in turn.

#### **SERVICES**

Guidance in a number of countries has put forth the concept of comprehensive wellness programmes. 'Comprehensiveness' means providing a range of offerings that target a number of health and wellbeing outcomes. Importantly, they need to be of interest to employees and as such the programmes need to be developed in consultation with employees [45, 46]. There are a number of challenges. Even when services meet the needs of certain types of employees, they may be inadequate or unattractive to other employees. One study found significant differences in the effect of workplace factors on health outcomes in male employees versus female employees [47]. The literature suggests that more components indicate a more effective wellness programme, both

in terms of participation and health and wellbeing outcomes. However, more is not always better. Recent research conducted by RAND Europe using Britain's Healthiest Workplace data found that simply offering more services without components that drive participation in, and the promotion of, health and wellbeing programmes is likely to be ineffective [48]. Services can also have more effect when combined. Some studies show that programmes combining both occupational health and safety (e.g. exposure to hazardous substances) with interventions on health behaviours (e.g. smoking) may prove more effective than programmes focused on either intervention in isolation [49].

#### **INCENTIVES**

Employees are only likely to benefit from health and wellbeing programmes if they participate. Participation remains a challenge for any health and wellbeing programme. Participation rates are typically below 50% of eligible employees [50, 51, 52]. In Healthiest Workplace by AIA Vitality, we see very low participation rates below 10% of eligible employees for a number of interventions that employers offer, which can undermine the logic for offering health and wellbeing programmes. Incentives are increasingly used to motivate participation. However, the level, type, and structure of incentives vary across health and wellbeing programmes, with some organisations using positive incentives (monetary or non-monetary) that may lead to higher participation, while others use alternative incentive structures like penalties for non-participation. Recent research in the US unsurprisingly suggests that incentives or penalties have different effects across sectors and worker groups (e.g. female or male workers; older and younger workers [53]). Finally, access is important. Employees clearly prefer programmes conducted at convenient times and places, including paid time off during the workday [46].

#### **PROMOTION**

Roughly half of all employers offer some type of health and wellbeing programme [54]. The data collected through Healthiest Workplace by AIA Vitality clearly show a significant gap between what employers say they offer and what employees perceive the offer to be. It appears that awareness of some interventions among employees is low and as such the take-up of interventions is low as well. Not surprisingly, employees who use the intervention report that the intervention has benefited their health and wellbeing. This speaks to the priorities of a programme aligning with the priorities of the participant. Communication clearly needs to go beyond raising awareness of what a health and wellbeing offer is and also allow the views of employees to be represented in the design and implementation of the health and wellbeing programme. Communication

can also improve the outcomes of a programme. One study showed that multiple follow-up messages throughout the running of the programme can be more successful in affecting behaviour and outcomes than one larger educational session at the start of a programme [55].

#### **LEADERSHIP**

Leadership is often recognised as one of the most important drivers of organisational change [56]. Having management involved in health promotion efforts has been linked with increased impact in terms of selfreported health and absenteeism [57]. Acknowledging the role of the workplace as a setting for health promotion is a view not necessarily held by all levels of management [52]. A qualitative study of three Canadian companies implementing interventions to manage musculoskeletal conditions found that while senior management was supportive of the interventions, middle-management was apathetic or, in one case, resistant to change [58]. In this case, participation in the programme remained low and as a result, the effectiveness of the programme was limited. The Britain's Healthiest Workplace data show that those organisations that discuss health and wellbeing at board level, report on health and wellbeing internally, or have engaged senior leadership have significantly higher participation in their health and wellbeing programmes than organisations that do not. The presence of these components by itself is also associated with higher participation in programmes.



#### **EVALUATION**

An essential component of a health and wellbeing programme is a continuous, systematic evaluation of design and impact. However, evaluations are rare among wellness programmes and where evaluation does exist, few are systematic or continuous. This is potentially a reason for the mixed results from wellness programme evaluations and the disconnect between best practices and strategies in use [59].

One possible reason for the disconnect between research and practice in wellness programmes may be the burden that regular evaluation of programme impact presents to employers, although it seems obvious that it is impossible to know what, and to what extent, a programme is effective without proper monitoring and evaluation in place. The experience from Healthiest Workplace by AIA Vitality is that organisations with better management information on health and wellbeing also tend to have higher participation in wellbeing programmes. In addition, employers are becoming increasingly interested in building the evidence base around health and wellbeing programmes.

# CONCLUSION

The health and wellbeing of employees makes an important contribution to business success. For this reason, a growing number of employers perceive employee wellbeing programmes as a strategic priority, particularly during challenging economic times.

Proactive management of employees' physical and mental health can produce a range of important business benefits including a reduction in sickness absence, improved productivity and staff engagement, lower staff turnover and a reduction in costs associated with recruitment.

This report provides you with an overview of the employees' unique clinical and lifestyle risk factors and assesses how they are responding to, and engaging in, the existing facilities and services offered.

Collectively this data provides a platform to implement employee wellbeing programmes to mitigate these risks and track organisation's progress. Even a small ongoing investment in the wellbeing of employees can pay big dividends for the business and help to improve the bottom line.


# APPENDICES

# APPENDIX A


#### **DEMOGRAPHIC BREAKDOWN OF RESPONDENTS**

The 2,137 employees that finished the survey in Sri Lanka are represented in the following graphs.


## DISTRIBUTION OF RESPONDENTS BY GENDER




### DISTRIBUTION OF RESPONDENTS BY AGE GROUPS



### DISTRIBUTION OF RESPONDENTS BY JOB CATEGORY



### DISTRIBUTION OF RESPONDENTS ACROSS INCOME GROUPS\*



<sup>\*</sup>Respondents who preferred not to provide answers are not included.

# APPENDIX B

### YEAR-ON-YEAR RESULTS\*

CATEGORY	INDICATOR	2017	2018	2019
Key outcomes				
Long-term health	AIA Vitality Age gap (in years)	-	5.2	4.9
Short-term work impairment	Days lost per employee per year	-	49.3	47.9
Work engagement	% low engaged	-	8.3%	10.3%
Determinants of he	ealth, productivity and engagement			
Lifestyle health	% with insufficient physical activity*	-	-	58.6%
	% not eating a healthy diet	-	91.5%	88.9%
	% current smokers	-	12.8%	13.6%
	% exceeding alcohol guidelines	-	1.1%	1.9%
	% sleeping less than 7 hours per night	-	46.4%	46.4%
Clinical health	% obese	-	10.6%	10.3%
	% with severe musculoskeletal problems**	-	7.6%	7.4%
Mental health	% with moderate or severe symptoms of depression	-	12.9%	15.0%
	% with 1 or more work-related stress factors	-	59.5%	56.1%
	% with financial concerns	-	33.6%	30.9%
Employee characte	eristics			
Survey respondent	% male	-	61.3%	59.1%
profile	Average age	-	32.5	32.0
	Average income	LKR.	LKR 92,595	LKR 104,761

<sup>\*</sup> The physical inactivity indicator in 2018 is not comparable to other years due to changes in the measurement method.

<sup>\*\*</sup> This question was not asked in Australia in 2018.


# APPENDIX C.

#### **CALCULATION OF PRODUCTIVITY LOSS**

The Work Productivity and Activity Impairment (WPAI) scale provides an estimate of employee absence based on absence over the week prior to the survey. This measures absence due to health problems and is based on employee self-reporting. Absence is estimated by the answer to the following question:

"During the past seven days, how many hours did you miss from work because of your health problems? Include hours you missed on sick days, times you went in late, left early, etc., because of your health problems. Do not include time you missed to participate in this study."

#### ABSENCE DUE TO HEALTH PROBLEMS




Overall, 82% of respondents in Sri Lanka did not miss any work time due to health problems.


To measure presenteeism - and therefore to get a sense of reduced productivity at work - the general health WPAI (WPAI-GH) scale asks six questions (including the question on absence above). Based on the numerical answers, a number of estimates can be made which assess productivity loss when multiplied with employee wages. In short, to capture presenteeism the WPAI scale takes the total number of hours an employee has actually worked during a week (taking absence and other loss of work into account) and multiplies this measure by the 'percent of self-estimated level of impact of health problems on productivity during the past seven days'.

Presenteeism is indicated by the answers to the following questions:

"During the past seven days, how many hours did you actually work?"



"During the past seven days, how much did your health problems affect your productivity while you were working?"



Overall, 43% of the respondents in Sri Lanka experienced no impact of health problems on productivity while working, compared to the benchmarks of 32% for the Asia Pacific and 43% for Sri Lanka's 2018 results.

# APPENDIX D

### RISK FACTOR DEFINITIONS USED IN THIS REPORT

Below is a description of the risk factors referred to in this report and what is considered to be the healthy range (and thus classified as low risk) for each:

HEALTH INDICATOR	DESCRIPTION	HEALTHY RANGE		
Body Mass Index (BMI)	BMI is a measure of a person's weight in relation to their height	Between 18.5 to 24.9		
Exercise	Cardiovascular exercise of moderate intensity, i.e. the person will breathe heavily, but should be able to talk.	At least 150 minutes of moderate intensity exercise per week or 75 minutes of vigorous exercise		
Smoking	Smoking status (current, former or non-smoker)	Non-smoker		
Nutrition	Fruit and vegetables	At least 5 servings per day		
Nutrition is a broad and	Whole grains	At least 3 servings per day		
complex risk factor. For the purpose of this	Lean meats	Choose lean meats		
survey, nutrition was	Low fat diary	Choose low fat or fat-free dairy		
assessed according to the following elements	Added fat	Avoid added fats		
the following elements	Trans fats	Avoid trans fats		
	Added salt	Avoid added salt		
	Sugary drinks	Less or equal to 1 drink per day		
Alcohol	Alcohol intake where one unit is equal to 8mg of alcohol	For men and women: no more than 14 units per week		
Mental wellbeing	Defined using the six-item Kessler Psychological Distress Scale	Low levels of psychological distress		
Screening	Participation in annual screening	All three health screenings (blood pressure, cholesterol and glucose) completed in the past year, regardless of the results of the checks		
Blood pressure	Blood pressure measured in mmHg	120/80 mmHg or less		
Cholesterol	Random total cholesterol level measured in mmol/l	Less than 200 mg/dL		
Glucose	Random total glucose level measured in mmol/l	Less than 140 mg/dL		

# **APPFNDIX F**

#### THE TEAM BEHIND THE HEALTHIEST WORKPLACE COMPETITION



#### ABOUT AIA & THE HEALTHIEST WORKPLACE BY AIA VITALITY

AIA Group Limited and its subsidiaries (collectively "AIA" or the "Group") comprise the largest independent publicly listed pan-Asian life insurance group. It has a presence in 18 markets in Asia-Pacific - wholly-owned branches and subsidiaries in Hong Kong, Thailand, Singapore, Malaysia, China, Korea, the Philippines, Australia, Indonesia, Taiwan, Vietnam, New Zealand, Macau, Brunei, Cambodia, a 97 per cent subsidiary in Sri Lanka, a 49 per cent joint venture in India and a representative office in Myanmar.

AIA meets the long-term savings and protection needs of individuals by offering a range of products and services including life insurance, accident and health insurance and savings plans. The Group also provides employee benefits, credit life and pension services to corporate clients. Through an extensive network of agents, partners and employees across Asia-Pacific, AIA serves the holders of 34 million individual policies and over 16 million participating members of group insurance schemes.

At AIA, we believe in the power of insurance to make a positive difference in people's lives. We do this through our products and services as well as by actively promoting the health and well-being of our customers, employees and agents and the communities in which they live and work. We are committed to helping people to live healthier, longer, better lives.

Investing in the health and wellbeing of employees can save businesses billions of dollars every year. For example, in Asia Pacific, an average of more than 67.5 days<sup>18</sup> are lost to absenteeism and presenteeism per employee every year, so identifying and addressing risks to employee health is critical to business performance.

For employees, participating in The Healthiest Workplace by AIA Vitality will give them a comprehensive overview of their health and wellbeing in terms of lifestyle, clinical indicators, mental health, stress and other areas that may concern them. Each participating employee will also receive a personal report detailing their unique Vitality Age.

Given that so much of our time is spent at work, we think it's important that we understand, measure and ultimately, improve wellbeing in the workplace. That's why we have created The Healthiest Workplace by AIA Vitality.

18

Source: The Healthiest Workplace by AIA Vitality 2019



#### **RAND EUROPE**

RAND Europe is an independent not-for-profit research institute whose mission is to help improve policy and decision-making through research and analysis. It realises its mission by undertaking objective, balanced, and relevant research and analysis; communicating findings to a wide audience, often through publications, many of which are available on its website; working in partnership with clients; and working collaboratively with others. RAND Europe's work lies on the continuum between that of universities and consultancies, combining the academic rigour of universities and the professional, task oriented approach of consultancies. It has uniquely broad experience in the fields of health and employment, having worked for a wide range of government and charitable funders globally. RAND Europe is part of the global RAND Corporation.

RAND Europe has formal links with the University of Cambridge, for instance through the Cambridge Centre for Health Services Research (CCHSR).

# REFERENCES

- [1] Institute for Health Metrics and Evaluation (2018). Findings from the Global Burden of Disease Study 2017. Seattle, WA: IHME.
- [2] World Health Organization (2010). Global Recommendations on Physical Activity for Health. Geneva: WHO. As of 10 September 2019: whqlibdoc.who.int/publications/2010/9789241599979\_eng.pdf?ua=1[1]
- [3] Centers for Disease Control and Prevention, Physical Activity and Health (2019). As of 10 September 2019: https://www.cdc.gov/physicalactivity/basics/pa-health/index.htm
- [4] Diet, nutrition and the prevention of chronic disease. Report of a joint FAO/WHO expert consultation, Geneva, World Health Organization, 2003 (WHO Technical Report Series, No. 916).
- [5] FAO (2017). What is a serving, 'Focus' webpage. As of 10 September 2019: http://www.fao.org/english/newsroom/focus/2003/fruitveg2.htm
- [6] FAO (1997). Carbohydrates in human nutrition. FAO Food and Nutrition Paper 66. As of 10 September 2019: http://www.fao.org/docrep/w8079e/w8079e00.htm
- [7] WHO (2016). Salt reduction Fact sheet. As of 10 September 2019: http://www.who.int/mediacentre/factsheets/fs393/en/
- [8] Health and Social Care Information Centre (2015) Statistics on Smoking: England, 2015. As of 10 September 2019: www.hscic.gov.uk/ catalogue/PUB17526/stat-smok-eng-2015-rep.pdf
- [9] WHO (2015). WHO global report on trends in tobacco smoking 2000-2025. As of 10 September 2019: http://www.who.int/tobacco/publications/surveillance/reportontrendstobaccosmoking/en
- [10] WHO (2019). WHO report on the global tobacco epidemic, 2019. Geneva: World Health Organisation; 2019: 1-19. As of September 2019: https://apps.who.int/iris/bitstream/hand le/10665/326043/9789241516204-eng.pdf?ua=1
- [11] WHO (2019). WHO report on the global tobacco epidemic, 2019. Geneva: World Health Organisation; 2019: 144-154. As of 11 September 2019: https://apps.who.int/iris/bitstream/hand le/10665/326043/9789241516204-eng.pdf?ua=1
- [12] Goodchild M., Nargis N., Tursan d'Espaignet E., Global economic cost of smoking-attributable diseases (2018). 27:58-64..

- [13] Weng W., Ali S. F., Leonardi-Bee J. S. (2012). Smoking and absence from work: Systematic review and meta-analysis of occupational studies, Addiction 108: 307-319.
- [14] di Castelnuovo A., Costanzo S., Bagnardi V., Donati M. B., Iacoviello L., de Gaetano G. (2006). Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34 prospective studies, Archives of Internal Medicine, Volume 166 (22): 2437-2445.
- [15] Roerecke M., Rehm J. (2013). Alcohol use disorders and mortality: a systematic review and meta-analysis, Addiction 108 (9): 1562-1578.
- [16] Rehm J., Baliunas D., Borges G. L., Graham K., Irving H., Kehoe T., Parry C. D., Patra J., Popova S., Poznyak V., Roerecke M., Room R., Samokhvalov A. V., Taylor B. (2010). The relation between different dimensions of alcohol consumption and burden of disease: an overview, Addiction Vol. 105(5): 814-843.
- [17] WHO (2010) Global strategy to reduce the harmful use of alcohol. As of 10 September 2019: http://apps.who.int/iris/bitstream/10665/44395/1/9789241599931\_eng.pdf?ua=1
- [18] European Food Information Council (2014). Behaviour Change Models and Strategies. As of 10 September 2019: https://www.eufic.org/en/healthy-living/article/motivating-behaviour-change
- [19] Ohayon, M. M. (2002). Epidemiology of insomnia: what we know and what we still need to learn. Sleep medicine reviews, 6(2), 97-111.
- [20] Rosekind, M. R., Gregory, K. B., Mallis, M. M., Brandt, S. L., Seal, B., & Lerner, D. (2010). The cost of poor sleep: workplace productivity loss and associated costs. Journal of Occupational and Environmental Medicine, 52(1), 91-98.
- [21] Edwards, J. A., Webster, S., Van Laar, D., & Easton, S. (2008). Psychometric analysis of the UK Health and Safety Executive's Management Standards work-related stress Indicator Tool. Work & Stress, 22(2), 96-107.
- [22] Kessler, R.C., Andrews, G., Colpe, L.J., Hiripi, E., Mroczek, D.K., Normand, S.-L.T., Walters, E.E., & Zaslavsky, A. (2002). Short screening scales to monitor population prevalences and trends in nonspecific psychological distress. Psychological Medicine, 32, 959-976.

- [23] Hassan, E., Austin, C., Celia, C., Disley, E., Hunt, P., Marjanovic, S., Shehabi, A., van Dijk, L. V., van Stolk C. (2009). Health and well-being at work in the United Kingdom. Rand Technical Report 758.
- [24] Vinokur, A. D., Price, R. H., & Caplan, R. D. (1996). Hard times and hurtful partners: how financial strain affects depression and relationship satisfaction of unemployed persons and their spouses. Journal of personality and social psychology, 71(1), 166.
- [25] Bianchini F, Kaaks R, Vainio H. (2002). Overweight, obesity, and cancer risk, Lancet Oncology, Vol 3(9): 1005-1008.
- [26] Bogers, R. P., Bemelmans, W. J., Hoogenveen, R. T., Boshuizen, H. C., Woodward, M., Knekt, P., ... & Thorpe, R. J. (2007). Association of overweight with increased risk of coronary heart disease partly independent of blood pressure and cholesterol levels: a meta-analysis of 21 cohort studies including more than 300,000 persons. Archives of internal medicine, 167(16), 1720-1728.
- [27] Larsson S. C., Wolk A. (2007). Overweight, obesity and risk of liver cancer: a meta-analysis of cohort studies, British Journal of Cancer Vol 97: 1005-1008.
- [28] Bilger, M., Finkelstein, E. A., Kruger, E., Tate, D. F., & Linnan, L. A. (2013). The effect of weight loss on health, productivity and medical expenditures among overweight employees. Medical care, 51(6), 471.
- [29] Huang, Y., Cai, X., Li, Y., Su, L., Mai, W., Wang, S., ... & Xu, D. (2014). Prehypertension and the risk of stroke A meta-analysis. Neurology, 82(13), 1153-1161.
- [30] World Health Organization, International Society of Hypertension Writing Group (2003). 2003 World Health Organization (WHO)/ International Society of Hypertension (ISH) statement on management of hypertension. Journal of hypertension, 21(11), 1983-1992.
- [31] CDC (2018). Work-Related Musculoskeletal Disorders & Ergonomics. As of 11 September 2019: https://www.cdc.gov/workplacehealthpromotion/health-strategies/musculoskeletal-disorders/index.html
- [32] March, L., Smith, E. U., Hoy, D. G., Cross, M. J., Sanchez-Riera, L., Blyth, F., ... & Woolf, A. D. (2014). Burden of disability due to musculoskeletal (MSK) disorders. Best practice & research Clinical rheumatology, 28(3), 353-366.

- [33] Koopmanschap, M., Burdorf, A., Lotters, F. (2013). Work Absenteeism and Productivity Loss at Work, In: Loisel P, Anema J (eds) Handbook of Work Disability: Prevention and Management, New York: Springer.
- [34] Brooks, A., Hagen, S. E., Sathyanarayanan, S., Schultz, A. B., Edington D. W. (2010). Presenteeism: critical issues. Journal Occupational Environmental Medicine 52(11):1055-67.
- [35] Mattke, S., Balakrishnan, A., Bergamo, G., Newberry, S. J. (2007). A review of methods to measure health-related productivity loss, American Journal Managed Care 13(4):211-7.
- [36] Hafner, M., van Stolk, C., Saunders, K., Krapels, J., Baruch, B. (2015). Health, wellbeing and productivity in the workplace: A Britain's Healthiest Company summary report, RAND Europe report. As of 11 September 2019: https://www.rand.org/pubs/research\_reports/RR1084.html
- [37] Harter, J. K., Schmidt, F. L., Hayes, T. L. (2002). Business-unit-level relationship between employee satisfaction, employee engagement, and business outcomes: A meta-analysis, Journal of Applied Psychology, Vol 87(2): 268-279.
- [38] Schaufeli, W. B., Salanova, M., González-Romá, V., & Bakker, A. B. (2002). The measurement of engagement and burnout: A two sample confirmatory factor analytic approach. Journal of Happiness studies, 3(1), 71-92.
- [39] Black, C. (2008). Working for a healthier tomorrow. Report for the Secretary of State for Health and the Secretary of State for Work and Pensions. As of 11 September 2019: https://www.gov.uk/government/uploads/system/uploads/attachment\_data/file/209782/hwwb-working-for-a-healthier-tomorrow.pdf
- [40] Hassan, E., Austin, C., Celia, C., Disley, E., Hunt, P., Marjanovic, S., Shehabi, A., van Dijk, L. V., van Stolk C. (2009). Health and well-being at work in the United Kingdom. Rand Technical Report 758.
- [41] Boorman, S. (2009). NHS Health and Well Being. Final Report. London: Department of Health. As of 11 September 2019: http://www.healthyregions.eu/uk/Boorman%20NHS%20HWB%20Final%20Report%20Nov%2009.pdf

- [42] Anderson, B. K., Larimer, M. E. (2002). Problem drinking and the workplace: an individualized approach to prevention. Psychology of Addictive Behaviors 16(3):243-251.
- [43] Osilla, K. C., Van Busum, K., Schnyer, C., Larkin, J. W., Eibner, C., Mattke, S. (2012). Systematic review of the impact of worksite wellness programs, American Journal of Managed Care, Vol.18, Issue 2: e68-e81.
- [44] Mattke, S., Liu, H., Caloyeras, J. P., Huang, C. Y., Van Busum, K. R., Khodyakov, D., Shier, V. (2013). Workplace Wellness Programs Study, Final Report. Santa Monica: RAND
- [45] Centers for Disease Control and Prevention. Essential Elements of Effective Workplace Programs and Policies for Improving Worker Health and Wellbeing. (2015). As of 11 September 2019: http://www.cdc.gov/niosh/twh/essentials.html
- [46] Consensus Statement of the Health Enhancement Research Organization. (2012). Guidance for a reasonably designed, employer-sponsored wellness program using outcomes-based incentives. Journal of Occupational and Environmental Medicine, 54(7), 889-896.
- [47] Peterson, M. (2004). What men and women value at work: Implications for workplace fhealth. Gender medicine, 1(2), 106-124.
- [48] Batorsky, B., Van Stolk, C., & Liu, H. (2016). Is More Always Better in Designing Workplace Wellness Programs?: A Comparison of Wellness Program Components Versus Outcomes. Journal of occupational and environmental medicine, 58(10), 987-993.
- [49] Barham, K., West, S., Trief, P., Morrow, C., Wade, M., Weinstock, R. S. (2011). Diabetes prevention and control in the workplace: a pilot project for county employees. Journal Public Health Management Practice 17(3):233-241.
- [50] Person, A. L., Colby, S. E., Bulova, J. A., & Eubanks, J. W. (2010) Barriers to participation in a worksite wellness program. Nutrition research and practice, 4(2), 149-154.
- [51] Robroek, S. J., Van Lenthe, F. J., Van Empelen, P., & Burdorf, A. (2009). Determinants of participation in worksite health promotion programmes: a systematic review. International Journal of Behavioral Nutrition and Physical Activity, 6(1), 26.

- [52] Linnan, L. A., Sorensen, G., Colditz, G., Klar, N., & Emmons, K. M. (2001). Using theory to understand the multiple determinants of low participation in worksite health promotion programs. Health education & behavior, 28(5), 591-607.
- [53] Batorsky, B., Taylor, E., Huang, C., Liu, H., & Mattke, S. (2016). Understanding the relationship between incentive design and participation in US workplace wellness programs. American Journal of Health Promotion, 30(3), 198-203.
- [54] Buck Consultants, L. L. C. (2012). Working well: A global survey of health promotion and workplace wellness strategies. Buck Consultants.
- [55] Heirich, M., Sieck, C. J. (2000). Worksite cardiovascular wellness programs as a route to substance abuse prevention. Journal Occupational Environmental Medicine 42(1):47-56.
- [56] Martins, E. C., & Terblanche, F. (2003). Building organisational culture that stimulates creativity and innovation. European journal of innovation management, 6(1), 64-74.
- [57] Dellve, L., Skagert, K., & Vilhelmsson, R. (2007). Leadership in workplace health promotion projects: 1-and 2-year effects on long-term work attendance. European Journal of Public Health, 17(5), 471-476.
- [58] Dixon, S., Theberge, N., & Cole, D. (2009). Sustaining Management Commitment to Workplace Health Programs: The Case of Participatory Ergonomics. Relations Industrielles/Industrial Relations, 64(1), 50-74.
- [59] Harden, A., Peersman, G., Oliver, S., Mauthner, M., & Oakley, A. (1999). A systematic review of the effectiveness of health promotion interventions in the workplace. Occupational medicine, 49(8), 540-548.